

MEDICARE FORM

Cimzia® (certolizumab pegol) Injectable Medication Precertification Request

Page 1 of 3

(All fields must be completed and legible for precertification review.)

For Medicare Advantage Part B: Phone: <u>1-866-503-0857</u> (TTY: <u>711</u>) FAX: <u>1-844-268-7263</u>

For other lines of business:

Please use other form.

Note: Cimzia is non-preferred.
Preferred products vary based on indication. See section G below.

Please indicate: Start of treatment: Start date				maication. O	se section o below.	
☐ Continuation of therapy: Date of	last treatment/					
Precertification Requested By:		Phone:		Fax: _		
A. PATIENT INFORMATION						
First Name:	Last Name:			DOB:		
Address:	City:			State:	ZIP:	
Home Phone: Work Phone:	(Cell Phone:		Email:		
Patient Current Weight: lbs or kgs Patie	nt Height: inches	or cms Alle	rgies:			
B. INSURANCE INFORMATION						
	Does patient have other coverage?		_			
	If yes, provide ID#: Carrier Name:		ier Name:			
	Insured:					
Medicare: ☐ Yes ☐ No If yes, provide ID #:	Med	icaid: 🗌 Yes 🗌 No	If yes, prov	ide ID #:		
C. PRESCRIBER INFORMATION	I. (N	(0)	/ O \ F			
	Last Name:	(Cr	еск Опе): _		. N.P. P.A.	
	City:		1	State:	ZIP:	
		NPI #:	DEA #:	1	UPIN:	
Provider Email:	Office Contact Name:			Phone:		
Specialty (Check one): Gastroenterologist Rhe	eumatologist 🔲 Derr	matologist	·			
Self-administered □ Physician's Office □ Home □ Outpatient Infusion Center Phone: □ Center Name: □ Home Infusion Center Phone: □ Administration Phone: □ Administration Center Phone: □ Administration Phone: <	ZIP:	Dispensing Provid Physician's Offic Specialty Pharm Other: Name: Address: City: Phone: TIN: NPI:	ce [Retail Pharn Mail Order State: Fax: PIN:	ZIP:	
Request is for Cimzia (certolizumab pegol)						
Dose: Frequence	cy:		CS Code:			
F. DIAGNOSIS INFORMATION - Please indicate primary IC						
Primary ICD Code:	_ Secondary ICD Code:			r ICD Code:		
Note: Cimzia is non-preferred. Entyvio, Inflectra, Remicade, Simponi Aria, and unbranded infliximab are preferred for MA plans. For MAPD plans, Enbrel, Humira, Kevzara, Otezla, Rinvoq, Skyrizi, Stelara, and Xeljanz/Xeljanz XR are preferred. Preferred products vary based on indication. Yes						



MEDICARE FORM

Cimzia® (certolizumab pegol) Injectable **Medication Precertification Request**

Page 2 of 3

(All fields must be completed and legible for precertification review.)

For Medicare Advantage Part B: Phone: <u>1-866-503-0857</u> (TTY: <u>711</u>) FAX: <u>1-844-268-7263</u>

For other lines of business:

Please use other form.

Note: Cimzia is non-preferred. Preferred products vary based on indication. See section G.

Patient First Name	Patient Last Name	Patient Phone	Patient DOB				
G. CLINICAL INFORMATION (continue	ed) – Required clinical information	n must he completed in its entirety fo	or all precertification requests				
			red products when indicated for the patient's				
diagnosis (select all that apply)							
 ☐ Enbrel (etanercept) ☐ Humira (adalimumab) ☐ Kevzara (sarilumab) ☐ Otezla (apremilast) ☐ Rinvoq (upadacitinib) ☐ Skyrizi (risankizumab-rzaa) ☐ Stelara (ustekinumab) ☐ Xeljanz/Xeljanz XR (tofacitinib) 							
GRYNZI (HSANRIZUMA		7) Meljaniz/Keljaniz XIX (tolacitimic)				
For All Requests (clinical documentati	on required for all requests):						
		other biologic (e.g., Humira) or targe	eted synthetic disease-modifying anti-rheumatic				
	umiant, Otezla, Xeljanz)? eived (including current utilizers)	a biologic (e.g., Humira) or targeted	synthetic DMARD (e.g., Olumiant, Xeljanz)				
	eased risk of tuberculosis?	test (e.a. tuherculosis skin test IPPI	D], interferon-release assay [IGRA], chest x-ray)	١			
	6 months of initiating therapy?	icat (c.g., tabercalosis skiri test [i i i	of, interioral release assay [refort], enest x-ray)	,			
		interferon-gamma assay (IGRA)					
		sis (TB) test:					
•	t tuberculosis 🔲 Yes 🔲 No 🕒	las treatment for latent tuberculosis	(TB) infection been initiated or completed?				
For Initiation Requests (clinical docum		Please select: treatment initiated	☐ treatment completed				
Ankylosing spondylitis and axial spor							
Please indicate loading dose at weeks (• •				
Please select which of the following app							
spondyloarthritis?	ceived (including current utilizers) a biologic (e.g., Humira) indicated	or active ankylosing spondylitis or active axial				
			nsteroidal anti-inflammatory drugs (NSAIDs), or				
does the Crohn's disease	he patient have an intolerance or	contraindication to at least two NSA	IDs?				
Please indicate loading dose at weeks 0	, 2, and 4: Please indica	ate maintenance dose:	frequency:weeks				
Yes No Has the patient been dia				-0			
	eived (including current utilizers) ne patient have fistulizing Crohn's		or moderately to severely active Crohn's disease	e?			
Yes No Has the patient tried and had an inadequate response to at least one conventional therapy option?							
Yes No Does the patient have a contraindication or intolerance to at least one conventional therapy option (e.g., azathioprine [Azasan, Imuran], budesonide [Entocort EC], ciprofloxacin							
		-	prednisolone [Solu-Medrol], methotrexate IM or \$	SC,			
	metronidazole [Flagyl], prednisone, sulfasalazine [Azulfidine, Sulfazine], rifaximin [Xifaxan],						
	tacrolim → Please select: ☐ Sulfas	us)? alazine (Azulfidine, Sulfazine)	letronidazole (Flagyl)				
☐ Ciprofloxacin (Cipro) ☐ Prednisone ☐ Budesonide (Entocort EC) ☐ Azathioprine (Azasan, Imuran)							
☐ Mercaptopurine (Purinethol)☐ Methotrexate IM or SC☐ Methylprednisolone (Solu-Medrol)☐ Rifaximin (Xifaxan)☐ Tacrolimus							
Plaque psoriasis							
Please indicate loading dose at weeks (· · · · · · · · · · · · · · · · · · ·		frequency:weeks				
☐ Yes ☐ No Has the patient been di ☐ Yes ☐ No Has the patient ever re	•		ndicated for the treatment of moderate to severe	re			
plaque psoriasis?		,					
		t, face, neck, scalp, genitals/groin, i	ntertriginous areas) affected? o starting the requested medication):%				
If less than 10% of BSA		surface area (BSA) affected (prior t	o starting the requested medication)/0				
			to phototherapy (e.g., UVB, PUVA) or				
	acologic treatment with methotre		ogic treatment with methotrexate, cyclosporine				
	and acitretin?	a cimical reason to avoia pharmase	ogic accument man meaner exact, cyclooperine				
		reason to avoid pharmacologic trea					
☐ Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease ☐ Breastfeeding ☐ Cannot be used due to risk of treatment-related toxicity ☐ Drug interaction							
	☐ Pregnancy or curre	ntly planning pregnancy					
	Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias,						
	uncontrolled hypert ☐ Other, please expla	•					



MEDICARE FORM

Cimzia® (certolizumab pegol) Injectable **Medication Precertification Request**

Page 3 of 3

(All fields must be completed and legible for precertification review.)

For Medicare Advantage Part B: Phone: <u>1-866-503-0857</u> (TTY: <u>711</u>) FAX: <u>1-844-268-7263</u>

For other lines of business: Please use other form.

Note: Cimzia is non-preferred. Preferred products vary based on indication. See section G.

Patient First Name	Patient Last Name	Patient Phone	Patient DOB					
G. CLINICAL INFORMATION (continu	red) – Reguired clinical information must	be completed in its entirety	for all precertification requests.					
Psoriatic arthritis								
	0, 2 and 4: Please indicate ma		frequency:weeks					
Yes No Has the patient been diagnosed with active psoriatic arthritis (PsA)?								
Yes No Does the patient have psoriatic arthritis with co-existent plaque psoriasis?								
Rheumatoid arthritis	0. 2 and 4: Diseas indicate ma	intononos doso:	fraguenavi					
	0, 2 and 4: Please indicate madiagnosed with moderately to severely ac							
	Yes No Has the patient ever received (including current utilizers) a biologic (e.g., Humira) or targeted synthetic disease modifying drug (DMARD) (e.g., Rinvoq, Xeljanz) indicated for moderately to severely active rheumatoid arthritis?							
└────────────────────────────────────	ne patient been tested for the rheumatoid	I factor (RF) biomarker?						
Please	e indicate test result: 🗌 positive 🛛 neg	gative not completed						
	ne patient been tested for the anti-cyclic		CP) biomarker?					
	e indicate test result: ☐ positive ☐ neg ne patient been tested for the C-reactive							
	e indicate test result: positive neg	· <u>`</u> ,						
	ne patient been tested for the erythrocyte		piomarker?					
Pleas	e indicate test result: positive neg	gative not completed						
		sponse after at least 3 mont	hs of treatment with methotrexate at a dose greater					
	or equal to 20mg per week?	. :	2					
	es No Has the patient experienced an es No Does the patient have a contra							
	Please indicate the contraind		•					
	☐ History of intolerance or a		pairment Hypersensitivity					
	☐ Clinical diagnosis of alcoh	ol use disorder, alcoholic liv	ver disease or other chronic liver disease					
	☐ Elevated liver transaminas	ses Significant drug inte	eraction Myelodysplasia Breastfeeding					
	☐ Interstitial pneumonitis or o	clinically significant pulmona	ary fibrosis					
	☐ Pregnancy or currently pla	• • • •						
	☐ Blood dyscrasias (e.g., thr							
For Continuation Bossests (clinical d	Other, please explain:		<u> </u>					
Please indicate maintenance dose:	ocumentation required for all requests	<u>5).</u>						
	receiving the requested drug through sar	mnlee or a manufacturer's r	vatient assistance program?					
			ase activity or improvement in signs and symptoms					
	t with the requested drug?		,					
Ankylosing spondylitis and axial spo	ndyloarthritis							
Please indicate which of the following has the patient experienced:								
☐ Functional status ☐ Total spinal pain ☐ Inflammation (e.g., morning stiffness) ☐ None of the above								
Crohn's disease								
Yes No Has the patient achieved or maintained remission?								
Please indicate which of the following h	nas the patient experienced: Abdominal mass	iarrhea	pearance of the mucosa					
	scoring tool (e.g., Crohn's Disease Activit							
Plaque psoriasis	3 (. 3 ,	,,						
	enced a reduction in body surface area (BSA) affected from baselin	e?					
Yes No Has the	e patient experienced an improvement in	signs and symptoms of the	e condition from baseline (e.g., itching, redness,					
-	, scaling, burning, cracking, pain)?							
Psoriatic arthritis only								
Please indicate which of the following h	· · · · · · · · · · · · · · · · · · ·	thesitis □ Skin and/or nait	involvement None of the above					
Number of swollen joints ☐ Number of tender joints ☐ Dactylitis ☐ Enthesitis ☐ Skin and/or nail involvement ☐ None of the above Rheumatoid arthritis								
	activity improvement from baseline in ter	nder joint count, swollen joi	nt count, pain, or disability:					
H. ACKNOWLEDGEMENT								
Request Completed By (Signature	Required):		Date: / /					
Any person who knowingly files a request for authorization of coverage of a medical procedure or service with the intent to injure, defraud or deceive any								
			the purpose of misleading, commits a fraudulent					
	ubjects such person to criminal and civ		.					

The plan may request additional information or clarification, if needed, to evaluate requests.